

506.72 877
169322
578
H
THE

JOURNAL

OF

THE LINNEAN SOCIETY.

ZOOLOGY.

VOL. X.



LONDON:

SOLD AT THE SOCIETY'S APARTMENTS, BURLINGTON HOUSE;

AND BY

LONGMANS, GREEN, READER, AND DYER,

AND

WILLIAMS AND NORGATE,

1870.

LIST OF PAPERS.

	Page
ANDREWS, W., Esq., M.R.I.A.	
On <i>Ziphius Sowerbiensis</i>	67
BAIRD, WILLIAM, M.D., F.R.S., &c.	
Contributions towards a Monograph of the Species of <i>Annelides</i> belonging to the <i>Amphinomacea</i> , with a List of the known Species, and a Description of several new Species (belonging to the group) contained in the National Collection of the British Museum. To which is appended a short Account of two hitherto nondescript Annulose Animals of a larval character. (Plates IV., V., and VI.)	215
Remarks on several Genera of <i>Annelides</i> , belonging to the Group Eunicea, with a notice of such Species as are contained in Collection of the British Museum, and a description of some others hitherto undescribed ..	341
BLACKWALL, JOHN, F.L.S.	
A List of Spiders captured by Professor E. PERCEVAL WRIGHT, M.D., in the province of Lucca, in Tuscany, in the summer of 1863, with characters of such Species as appear to be new or little known to Arachnologists. (Plates XV. & XVI.) ..	405
CAMBRIDGE, Rev. O. P.	
Descriptions of a New Genus and Six New Species of Spiders. (Plate IX.)	264
Catalogue of a Collection of Ceylon Araneidea lately received from Mr. J. Nietner, with Descriptions of new Species and Characters of a new Genus.—Part I. (Plates XI., XII., XIII.)	373
Descriptions and Sketches of two new Species of Araneidea, with Characters of a new Genus. (Plate XIV.)	398
EDWARD, THOMAS, A.L.S.	
A few Additional Particulars regarding <i>Couchia Edwardii</i>	1
FOLIN, Marquis LÉOPOLD DE.	
Observations on the Septum of the Cæcidæ ; and some remarks on the subject of the suppression of the Genera <i>Brochina</i> and <i>Strebloceras</i> or <i>Phleboceras</i> . (Plate VIII.)	254

	Page
GREEN, ASHDOWN H., Esq.	
On the Natural History and Hunting of the Beaver (<i>Castor canadensis</i> , Kuhl) on the Pacific Slope of the Rocky Mountains. With Supplementary Notes by ROBERT BROWN, Esq., F.R.G.S.	361
KIRBY, W. F., Memb. Ent. Soc. Lond.	
On the necessity of a Reform in the Generic Nomenclature of Diurnal Lepidoptera, illustrated by a Review of the Genera proposed from the time of Linnæus to the year 1816	494
LUBBOCK, Sir JOHN, Bart., V.P.L.S., F.R.S., &c.	
Note on the Discovery of <i>Planaria terrestris</i> in England	193
McINTOSH, W. C., M.D., F.L.S.	
Note on the Development of Lost Parts in the <i>Nemertean</i> s. (Plate VII.)	251
M'LACHLAN, ROBERT, Esq., F.L.S., Sec. Ent. Soc.	
On some new Forms of Trichopterous Insects from New Zealand; with a List of the Species known to inhabit those Colonies. (Plate II.)	196
MURRAY, ANDREW, Esq., F.L.S.	
On an undescribed Light-giving Coleopterous Larva (pro- visionally named <i>Astraptor illuminator</i>). (Plate I.)	74
PASCOE, FRANCIS P., Esq., F.L.S., F.Z.S., &c.	
Contributions towards a Knowledge of the <i>Curculionidæ</i> .— Part I. (Plates XVII., XVIII. & XIX.)	434
SAUNDERS, EDWARD, Esq., F.L.S.	
Descriptions of ten new Species of the Genus <i>Paracupta</i> , H. Deyrolle, and of ten new Species of the Genus <i>Conognatha</i> , Escholtze. (Plate X.)	331
THEOBALD, W., Jun., Geological Survey of India.	
Catalogue of the Reptiles of British Birma, embracing the Provinces of Pegu, Martaban, and Tenasserim; with de- scriptions of new or little-known species.	4
TRIMEN, ROLAND, Memb. Ent. Soc. Lond.	
On the occurrence of <i>Astraptor illuminator</i> , Murray, or a closely allied insect, near Buenos Ayres	503
WALKER, FRANCIS, Esq., F.L.S.	
Catalogue of the Homopterous Insects collected in the Indian Archipelago by Mr. A. R. WALLACE, with descriptions of New Species. (Plate III.)	82, 276
WILSON, C. A., Esq., F.L.S.	
Note on <i>Moloch horridus</i>	69

Thlaosoma dubium.

- Fig. 25. Relative position of eyes.
 Fig. 26. Perspective view of fore part of caput, falces, maxillæ, and labium.
 Fig. 27. Ditto of cephalothorax.
 Fig. 28. Cephalothorax and abdomen from above and behind.
 Fig. 29. Profile of cephalothorax and falces.
 Fig. 30. Cephalothorax from above and behind, a more enlarged view.
 Fig. 31. Leg of first pair.
 Fig. 32. Underside, showing fore part of caput, falces, maxillæ, and labium.
 Fig. 33. Palpi and sternum, natural size.
 Fig. 34. Profile of Spider, with legs removed, from a different specimen.
 Fig. 35. Natural length.

T. distinctum.

- Fig. 36. View of Spider, from above.
 Fig. 37. Profile of cephalothorax and abdomen.
 Fig. 38. Natural length.

Catalogue of the Homopterous Insects collected in the Indian Archipelago by Mr. A. R. WALLACE, with descriptions of New Species. By FRANCIS WALKER, Esq., F.L.S.

(Continued from p. 193.)

Fam. CICADELLINA, *Burm.*Tribe LÆVIPEDES, *Amyot et Serv.*Subtribe CERCOPIDES, *Amyot et Serv.*Gen. PHILAGRA, *Stål.*

287. PHILAGRA DOUGLASI, *Stål*, *Trans. Ent. Soc. Lond.* 3rd ser. i. 593.

Hab. Batchian.

288. PHILAGRA SCOTTI, *Stål*, *Trans. Ent. Soc. Lond.* 3rd ser. i. 594.

Hab. Batchian.

Gen. CERCOPIS, *Fabr.*

289. CERCOPIS SPECTABILIS, *Burm.* *Nov. Act. Phys. Med. Nat. Cur.* xvi. *Suppl.* 304, pl. 41. f. 8.

Hab. Sumatra; inhabits also the Philippine Isles.

290. CERCOPIS PLANA, *Walk.* *Cat. Hom.* iii. 653.

Hab. Penang; inhabits also Java.

291. CERCOPIS UNDULIFERA. *Fæm.* *Nigra*, capite vitta incisa tes-

Male. Testaceous, stout. Head, thorax, and fore wings speckled with brown. Head conical, as broad as the thorax; vertex very much broader than long; front and face flat, piceous, the former bordered with pale yellow on each side. Pectus with two piceous stripes. Legs piceous. Fore wings with blackish speckles; a broad irregular blackish band extending from the base to full two-thirds of the length. Hind wings cinereous, hyaline; veins black. Length of the body 2 lines, of the wings 4 lines.

Hab. New Guinea.

Order PHYTOPHTHIRE, *Burm.*

Fam. PSYLLIDÆ, *Latr.*

Gen. CARSIDARA.

Mas. Corpus sat robustum. Caput thorace angustius, supra excavatum. Antennæ gracillimæ, corpore breviores; articuli 1^{us} et 2^{us} incrassati. Pedes breves, robusti. Alæ anticæ corpore multo longiores, vena longitudinali ramos quatuor emittente apicem versus arcuata, ramis 1°, 2°, 3°que furcatis. Alæ posticæ venis duabus longitudinalibus gracillimis.

Male. Body moderately stout. Head narrower than the thorax, depressed above. Antennæ very slender, shorter than the body; first and second joints incrassated; third, fourth and fifth successively shorter; sixth and seventh as long as the fifth; eighth longer than the seventh. Legs short, stout. Fore wings much longer than the body, with a vein which emits four branches and is curved at three-fourths of the length to the hind border near the tip; first branch emerging near the base, forked, its fore fork short, joining the stigma, its hind fork long, joining the costa beyond the stigma; second branch emerging very near the base, joining the hind border at much before the middle; third emerging beyond the first, joining the hind border close to the second, emitting near its tip a short curved fork to the hind border; fourth straight, emerging opposite the stigma, proceeding obliquely to the hind border. Hind wings with two very slender longitudinal veins.

442. CARSIDARA MARGINALIS. *Mas.* Testacea, capite lituris duabus lateralibus ferrugineis, antennis albidis articulis 1° et 2° ferrugineis, 3° et sequentibus apice nigris, prothorace guttis duabus lateralibus ferrugineis, mesothorace ferrugineo vittis quinque (quatuor interruptis)

pallide flavis, abdomine pedibusque fulvis, tibiis pallide flavis, alis diaphanis testaceo venosis, alis anticis quatuor punctis fuscis stigmatibus albedo basi apiceque nigro.

Male. Testaceous. Head with a ferruginous mark on each side between the eyes. Antennæ whitish; first and second joints ferruginous; all the following joints black towards the tips. Prothorax with a ferruginous dot on each side. Mesothorax ferruginous, with five pale yellow stripes, of which the middle one only is entire. Abdomen tawny, pale yellow beneath. Legs tawny; tibiæ pale yellow. Wings pellucid; veins testaceous. Fore wings with a whitish stigma, which is blackish at each end; four brown dots along the hind border. Length of the body $1\frac{3}{4}$ line, of the wings 5 lines.

Hab. Celebes.

Genus TYORA.

Mas. Corpus gracile. Antennæ graciles, filiformes, thorace multo longiores; articuli 1^{us} et 2^{us} incrassati. Pedes robusti, breviusculi. Alæ anticæ angustæ, venis tribus longitudinalibus basi connexis, 1^a vittas duas ad costam emittente, 2^a vittas tres ad marginem posticum emittente.

Male. Body slender. Antennæ slender, filiform, much longer than the thorax; first and second joints incrassated. Legs stout rather short. Fore wings narrow, with three longitudinal veins which are connected near the base; first vein emitting an oblique vein and an exterior short directly transverse vein to the costa; second vein emitting three oblique veins to the hind border (of these secondary veins the first and second are connected by a transverse vein, third vein extending obliquely to the hind border).

443. TYORA CONGRUA. *Mas.* Fulva, capite subtus pallide flavescente, thorace flavescente sexvittato, alis hyalinis subcinereis fulvo venosis.

Male. Tawny. Head pale yellowish beneath. Thorax with six pale yellowish stripes, the two exterior stripes broader than the others. Wings hyaline, slightly cinereous; veins tawny. Length of the body $1\frac{3}{4}$ line, of the wings 4 lines.

Hab. Mysol.

Fam. COCCIDÆ, Latr.

Gen. MONOPHLEBUS, Leach.

444. MONOPHLEBUS ATRIPENNIS, Burm. Handb. ii. 80.

Hab. Batchian; Celebes; Gilolo. Inhabits also Hindostan.

