

21. *Ephoria arenosa* Btlr. ♀
 22. " " f. *insularis* (nov.) ♂
 23. *Boarmia gorbatschevi* (sp. nov.) ♂
 24. *Itame fulvaria* Vill. *imperatoria* (subsp. nov.) ♂
 25. *Chiasma clathrata* L. *djakonovi* (subsp. nov.) ♂
 26. " " " " " " ♀
 27. *Aspilates formosaria* Ev. ab. *brunneata* (nov.) ♀
 Textfigur 1. *Abraxas grossulariata* L. Sexual-Armatur ♂
 " 2. *Abraxas heringi* m., Sexual-Armatur ♂.

Fauna sumatrensis.

(Beitrag Nr. 61).

Psyllidae.

By David L. Crawford, University of Hawai (Honolulu).

(with 1 Textfigure)

In a collection of *Psyllidae* (*Chermidae* of some authors) sent to me by Dr. E. Jacobson from Sumatra there are several previously known species represented and one new species which is of particular interest because of its possible relationship with a species which has become established in Hawaii.

Pauropsylla udei Rübsaamen¹): This easily recognized species, common throughout the old world tropics as gall-makers on *Ficus*, is represented by 6 specimens collected at Fort de Kock, by E. Jacobson, at 920 m elevation.

Paurocephala psylloptera Crawford²): This widely distributed species is represented by 15 specimens collected at Fort de Kock, Sumatra, by E. Jacobson on *Ficus fistulosa* at 920 m elevation. Most of the specimens have the dorsal thorax brown with orange streaks and spots.

Macrohormotoma gladiatum Kuwayama³): Six specimens of this species are in the collection, taken at Fort de Kock, Sumatra on *Ficus retusa* L. by E. Jacobson, some in February 1923 and one in 1924, at 920 m elevation.

This species was first located in Formosa, but subsequently has appeared in collections from Macao and the island of Larat.

Tenaphalara malayensis Crawford¹): This Malay Peninsula species is also in Sumatra and Java, for two specimens are very similar to the Singapore specimens on which the species description was based. One of

¹) Ent. Nachrichten Bd. XXV, p. 264, 1899 — Crawford, Philippine Journ. Science, Vol. 15, p. 145, 1919.


²) Philippine Journ. Science, Vol. 8, p. 294, 1913.

³) Trans. Sapporo Nat. Hist. Soc. Vol. II, p. 179, 1907. — Crawford, Proc. Haw. Ent. Soc. Vol. VI, Nr. 1, pp. 36—39, 1925.

these specimens is from Fort de Kock, taken by E. Jacobson in December 1920, at an elevation of 920 m. The other specimen is from Samarang, Java, taken by E. Jacobson in 1910.

Megatrioza mallotica (n. sp.): A large species, body (to tip of folded wings) from 6 to 8 mm long; body 3 to 4 mm long; forewing 5.7 mm long. General color light ochreous brown, sometimes with slight darker spots or streaks on thoracic dorsum; eyes and tips of antennae black.

Head somewhat declivous, with vertex about half as long as broad between eyes, clothed with long hairs; genal cones about half as long as vertex, small, subacute at apex; antennae about one and a half times



Megatrioza mallotica n. sp.

as long as width of head (including eyes). Thorax moderately broad, wings arched, with an abundance of long hairs on dorsal surface. Legs long and hairy; hind tibiae with small spur at base. Fore

wings very long, acutely pointed, flavous in color; venation as shown in illustration; hind wings less than half as long as forewings, clear. Abdomen: Female genital segment moderately long, about half as long as width of forewing. Male anal valve with lateral wings long; forceps long and slender.

Fort de Kock, Sumatra (E. Jacobson), 3 females and 2 males, collected December 1920 at 920 m elevation. This species forms galls on *Mallotus philippinensis* (Euphorbiaceae).

In several respects this species resembles its Hawaiian relative *M. palmicola* Crawford, but is quite distinct, however.

Hyménoptères Sud-Américains du Deutsches Entomologisches Institut: *Terebrantia*.

Par le Dr. Jean Brèthes.

(Supplement.)

[In den Ent. Mitteil. XVI, 1927, p. 296—309 und 319—335, ist eine Arbeit von dem leider inzwischen verstorbenen Dr. J. Brèthes über *Terebrantia* publiziert, bei welcher leider auf p. 309 der Schlußteil der Beschreibung von *Distantella Fischeri* (n. sp.), ebenso wie der der Tribus *Mesostenini* behandelnde Teil durch ein Versehen des Setzers au


¹⁾ Philippine Journ. Sc. Vol. 15, p. 165, 1919.

Band XVII, 1928

Entomologische Mitteilungen

(Organ der Wanderversammlungen Deutscher Entomologen)

Herausgegeben und redigiert von Walther Horn



All men of science are brothers. ...
Edgew. David August 1914)

Erschienen in 6 Einzelnummern
mit VI und 458 Seiten, 9 Tafeln und 172 Textfiguren.

| | | |
|-------|------------|----------------|
| Nr. 1 | erschienen | 5. Jan. 1928 |
| " 2 | " | 8. März 1928 |
| " 3 | " | 5. Mai 1928 |
| " 4 | " | 5. Juli 1928 |
| " 5 | " | 20. Sept. 1928 |
| " 6 | " | 20. Dez. 1928 |

Berlin-Dahlem
Goßler-Str 20.

Preis im Buchhandel 12,— M.

| | Seite |
|--|----------|
| Crawford, David L., <i>Psyllidae</i> (Hom.) (Mit 1 Fig.) | 425 |
| Malloch, J. R., Family <i>Muscidae</i> (Dipt.) (with 3 figures) | 290, 310 |
| Uhmann, E., <i>Hispinae</i> (Col.) (Mit 4 Fig.) | 370, 393 |
| Wasmann, E., <i>Paussidae</i> : Genus <i>Paussus</i> L. (Col.) (Mit 3 Photographien) | 238 |
| Werner, F., <i>Mantodea</i> (Orth.) | 412 |

Spezielle Systematik.

Orthoptera, Odonata.

| | |
|---|-----|
| Crampton, G. C., Anatomical evidence that <i>Cylindracheta</i> is a Gryllotalpoid not an Embiid. (Mit Taf. 4) | 252 |
| Fudakowski, Józef, Über eine melanotische Aberration von <i>Enallagma cyathigerum</i> Charp. (Odon.) (Mit 3 Textfig.) | 174 |
| Günther, Klaus, Beitrag zur Orthopterenfauna der Mark Brandenburg | 285 |
| Karny, H. H., Vorläufige Mitteilung über die wissenschaftlichen Ergebnisse meines Europa-Urlaubs | 60 |
| Krauss, Dr. H. A., Die Dermapteren und Orthopteren der balearischen Inseln. Mit 3 Textfig.) | 140 |
| Ris, Dr. F., Zwei neue Odonaten aus Chile und der argentinischen Kordillere Mit 12 Textfig.) | 162 |
| — Ein neuer <i>Gomphus</i> aus Shantung, China (Odon.) (Mit 4 Textfig.) | 273 |
| — <i>Enallagma deserti</i> Selys, eine vergessene Libelle. (Mit 2 Textfig.) | 277 |
| Schmidt, Erich, Bemerkungen über Lestiden (Odon.) (Mit 6 Textfig.) | 244 |
| — Zur Libellenfauna der Mark Brandenburg | 375 |
| Uvarow, B. P., A new <i>Tmethis</i> from S. W. Anatolia (Acridid.) | 176 |
| Werner, F., Fauna sumatrensis Nr. 60, <i>Mantodea</i> | 412 |

Hemiptera, Copeognatha.

| | |
|--|-----|
| Borgmeier, Th., Ein interessantes Copeognathen-Gespinnst aus Brasilien. (Mit 1 Fig.) | 155 |
| Crawford, David L., Fauna sumatrensis Nr. 61. <i>Psyllidae</i> (Mit 1 Fig.) | 425 |
| Nevsky, V. P., The plant-lice of Middle-Asia. I. (with plate 3) | 182 |

Coleoptera.

| | |
|--|----------|
| Benick, L., Alte und neue Steninen aus Australien und Ozeanien (Staph.) (Mit 3 Textfig.) | 177 |
| Bokor, Elemér, Bestimmungstabelle der Bathyscinen-Gattung <i>Sophrichaeta</i> Rtt. (Silph.) | 114 |
| Borchmann, F., Fauna sumatrensis Nr. 59 <i>Alleculidae</i> | 407 |
| Cameron, Malcolm, Fauna sumatrensis Nr. 54 <i>Staphylinidae</i> | 90 |
| Carter, St. J., Some new Australian <i>Heteromera</i> in the Germ. Entom. Mus. | 132 |
| Hajóss, Jos., Ein neuer <i>Centorrhynchus</i> aus Ungarn. (Curcul.) | 145 |
| Jedlička, A., Neue palaarktische Carabiden | 44 |
| Kleine, R., Neue <i>Metriorrhynchus</i> von Australien und Batjan (Lycid.) | 120 |
| Korschevsky, R., Bemerkungen über exotische <i>Coccinellidae</i> der alten Welt mit Beschreibung einer neuen Art. (Mit 8 Textfig.) | 41 |
| Nagel, P., Neues über Hirschkäfer (Lucan.) (Mit 3 Textfig.) | 257 |
| Théry, A., Buprestides nouveaux du Deutsches Entomologisches Museum. (2ème note) (Avec 2 figures) | 76 |
| Uhmann, E., <i>Probaenia crenatula</i> (n. sp.) nebst Fraßbild und <i>Alurnus Lydiae</i> (n. sp.) | 287 |
| — Fauna sumatrensis Nr. 57: <i>Hispinae</i> (Mit 4 Fig.) | 370, 393 |
| Wasmann, E., Fauna sumatrensis Nr. 55: <i>Paussidae</i> : Genus <i>Paussus</i> L. (Mit 3 Photographien) | 238 |